

订购热线: 4008-898-798

## Anti-F13A1 antibody

**Cat. No.** ml121318

**Package** 25 μl/100 μl/200 μl

**Storage** -20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol

**Product overview** 

**Description** Anti-F13A1 rabbit polyclonal antibody

Applications ELISA, IHC

**Immunogen** Fusion protein of human F13A1

ReactivityHuman, MouseContent0.1 mg/mlHost speciesRabbit

Ig classImmunogen-specific rabbit IgGPurificationAntigen affinity purification

**Target information** 

Symbol F13A1

Full name Coagulation factor XIII, A1 polypeptide

Synonyms F13A Swissprot P00488

## **Target Background**

This gene encodes the coagulation factor XIII A subunit. Coagulation factor XIII is the last zymogen to become activated in the blood coagulation cascade. Plasma factor XIII is a heterotetramer composed of 2 A subunits and 2 B subunits. The A subunits have catalytic function, and the B subunits do not have enzymatic activity and may serve as plasma carrier molecules. Platelet factor XIII is comprised only of 2 A subunits, which are identical to those of plasma origin. Upon cleavage of the activation peptide by thrombin and in the presence of calcium ion, the plasma factor XIII dissociates its B subunits and yields the same active enzyme, factor XIIIa, as platelet factor XIII. This enzyme acts as a transglutaminase to catalyze the formation of gamma-glutamyl-epsilon-lysine crosslinking between fibrin molecules, thus stabilizing the fibrin clot.



订购热线: 4008-898-798


## **Applications**

## **Immunohistochemistry**

Predicted cell location: Secreted, Cytoplasm Positive control: Human esophagus cancer

Recommended dilution: 25-100





The image on the left is immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using ml121318(F13A1 Antibody) at dilution 1/25, on the right is treated with fusion protein. (Original magnification: ×200)

**ELISA** 

Recommended dilution: 2000-5000


联系电话: 4008-898-798, 021-61725725


联系QQ: 2881505695, 2881505696

邮箱: mlbio\_cn@yeah.net 网址: www.mlbio.cn

Predicted cell location: Secreted, Cytoplasm

Positive control: Human lung cancer Recommended dilution: 25-100





The image on the left is immunohistochemistry of paraffin-embedded Human lung cancer tissue using ml121318(F13A1 Antibody) at dilution 1/25, on the right is treated with fusion protein. (Original

magnification: ×200)