

## Anti-CLEC4D antibody

|                 |                                                         |
|-----------------|---------------------------------------------------------|
| <b>Cat. No.</b> | ml120434                                                |
| <b>Package</b>  | 25 µl/100 µl/200 µl                                     |
| <b>Storage</b>  | -20°C, pH7.4 PBS, 0.05% NaN <sub>3</sub> , 40% Glycerol |

### Product overview

|                     |                                        |
|---------------------|----------------------------------------|
| <b>Description</b>  | Anti-CLEC4D rabbit polyclonal antibody |
| <b>Applications</b> | ELISA, WB                              |
| <b>Immunogen</b>    | Fusion protein of human CLEC4D         |
| <b>Reactivity</b>   | Human                                  |
| <b>Content</b>      | 0.6 mg/ml                              |
| <b>Host species</b> | Rabbit                                 |
| <b>Ig class</b>     | Immunogen-specific rabbit IgG          |
| <b>Purification</b> | Antigen affinity purification          |

### Target information

|                  |                                         |
|------------------|-----------------------------------------|
| <b>Symbol</b>    | CLEC4D                                  |
| <b>Full name</b> | C-type lectin domain family 4, member D |
| <b>Synonyms</b>  | MCL, MPCL, CLEC6, CLEC-6, CLECSF8       |
| <b>Swissprot</b> | Q8WXI8                                  |

### Target Background

This gene encodes a member of the C-type lectin/C-type lectin-like domain (CTL/CTLD) superfamily. Members of this family share a common protein fold and have diverse functions, such as cell adhesion, cell-cell signalling, glycoprotein turnover, and roles in inflammation and immune response. This gene is closely linked to other CTL/CTLD superfamily members on chromosome 12p13 in the natural killer gene complex region.

订购热线: 4008-898-798

#### Applications

##### Western blotting

Predicted band size: 25 kDa

Positive control: 293T cells

Recommended dilution: 500-2000

Gel: 10% SDS-PAGE


Lysate: 40 µg

Lane: 293T cells

Primary antibody: ml120434 (CLEC4D Antibody) at dilution 1/123

Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution

Exposure time: 1 minute



##### ELISA

Recommended dilution: 2000-5000

联系电话: 4008-898-798, 021-61725725

联系QQ: 2881505695, 2881505696

邮箱: [mlbio\\_cn@yeah.net](mailto:mlbio_cn@yeah.net)

网址: [www.mlbio.cn](http://www.mlbio.cn)